

Global well-posedness for the defocusing, quintic nonlinear Schrödinger equation in one dimension

Ben Dodson

October 20, 2009

Abstract: In this paper, we prove global well-posedness for low regularity data for the one dimensional quintic defocusing nonlinear Schrödinger equation. We show that a unique solution exists for $u_0 \in H^s(\mathbf{R})$, $s > \frac{8}{29}$. This improves the result in [13], which proved global well-posedness for $s > \frac{1}{3}$. The main new argument is that we obtain almost Morawetz estimates with improved error.

1 Introduction

In this paper we study the initial value problem for the quintic, defocusing, nonlinear Schrödinger equation in one dimension,

$$\begin{aligned} iu_t + \Delta u &= |u|^4 u, \\ u(0, x) &= u_0 \in H^s(\mathbf{R}). \end{aligned} \tag{1.1}$$

This is an L^2 -critical equation. By the results of [4], this equation has a local solution on some $[0, T]$, $T(\|u_0\|_{H^s(\mathbf{R})}) > 0$, when $s > 0$. If a solution to (1.1) fails to be global and only exists on $[0, T_*)$, $T_* < \infty$, then

$$\lim_{t \nearrow T_*} \|u(t)\|_{H^s(\mathbf{R})} = \infty. \tag{1.2}$$

[3] proved

$$M(u(t)) = \int |u(t, x)|^2 dx = M(u(0)), \tag{1.3}$$

$$E(u(t)) = \frac{1}{2} \int |\nabla u(t, x)|^2 dx + \frac{1}{6} \int |u(t, x)|^6 dx = E(u(0)), \tag{1.4}$$

are conserved, giving global well-posedness for $u_0 \in H^1(\mathbf{R})$. The regularity necessary for global well-posedness has since been lowered to $s > 1/3$, (see [13]). In this paper we will prove

Theorem 1.1 *(1.1) is globally well-posed for all $u_0 \in H^s(\mathbf{R})$, $s > \frac{8}{29}$. Moreover,*

$$\sup_{t \in [0, T]} \|u(t)\|_{H^s(\mathbf{R})} \lesssim (1 + T)^{\frac{4(1-s)s}{29s-8}+}. \quad (1.5)$$

[13] used the I-method, a method that we will utilize in this paper as well. The I-method was first introduced for the defocusing, cubic initial value problem (see [7]).

In §2, we will start with some preliminary information, including the Strichartz estimates, Littlewood-Paley theory, a description of the I-method, and a local well-posedness result. In §3, an energy increment will be obtained. In §4, the almost Morawetz estimates will be proved. In §5, we will prove the theorem.

2 Local Well-posedness

The proof of local well-posedness makes use of the Strichartz estimates.

Theorem 2.1 *A pair (p, q) is called an admissible pair if $\frac{2}{p} + \frac{1}{q} = \frac{1}{2}$. If (p, q) and $(\tilde{p}, \tilde{q})$ are admissible pairs and $u(t, x)$ solves*

$$\begin{aligned} iu_t + \Delta u &= F(t), \\ u(0, x) &= u_0, \end{aligned} \quad (2.1)$$

then

$$\|u(t, x)\|_{L_t^p L_x^q(J \times \mathbf{R})} \lesssim \|u_0\|_{L^2(\mathbf{R})} + \|F(t)\|_{L_t^{\tilde{p}'} L_x^{\tilde{q}'}(J \times \mathbf{R})}. \quad (2.2)$$

Proof: See [23]. p' denotes the Lebesgue exponent $\frac{p}{p-1}$.

The Strichartz space will be defined by the norm

$$\|u\|_{S^0(J \times \mathbf{R})} = \sup_{(p, q) \text{ admissible}} \|u\|_{L_t^p L_x^q(J \times \mathbf{R})}. \quad (2.3)$$

The space $N^0(J \times \mathbf{R})$ is the dual space to $S^0(J \times \mathbf{R})$. See [23] for more details.

We will also make use of the Littlewood-Paley decomposition. Suppose $\phi(x)$ is a smooth function,

$$\phi(x) = \begin{cases} 1, & |x| \leq 1/2; \\ 0, & |x| > 1. \end{cases} \quad (2.4)$$

$$\begin{aligned} \mathcal{F}(P_{\leq N}u) &= \phi\left(\frac{\xi}{N}\right)\hat{u}(\xi), \\ \mathcal{F}(P_{>N}u) &= (1 - \phi\left(\frac{\xi}{N}\right))\hat{u}(\xi), \\ \mathcal{F}(P_Nu) &= P_{\leq N}u - P_{\leq \frac{N}{2}}u. \end{aligned} \quad (2.5)$$

For convenience, let $u_N = P_Nu$, similarly for $u_{\leq N}$, $u_{>N}$.

The I - operator is a Fourier multiplier,

$$I_N : H^s(\mathbf{R}) \rightarrow H^1(\mathbf{R}), \quad (2.6)$$

$$\widehat{I_N f}(\xi) = m_N(\xi)\hat{f}(\xi), \quad (2.7)$$

$$m_N(\xi) = \begin{cases} 1, & |\xi| \leq N; \\ (\frac{N}{|\xi|})^{s-1}, & |\xi| > 2N. \end{cases} \quad (2.8)$$

$$\begin{aligned} \|Iu\|_{H^1(\mathbf{R})} &\lesssim N^{1-s}\|u\|_{H^s(\mathbf{R})}, \\ \|u\|_{H^s(\mathbf{R})} &\lesssim \|Iu\|_{H^1(\mathbf{R})}, \end{aligned} \quad (2.9)$$

therefore, controlling $E(Iu(t))$ gives control of $\|u(t)\|_{H^s(\mathbf{R})}$. For the rest of the paper, If denotes $I_N f$, and the presence of an N is implied.

Lemma 2.2 *Let I be a compact time interval, $t_0 \in I$, $N > 0$, and suppose u_1, u_2 are two solutions to (2.2) such that $u_j(t)$ has Fourier support in the region $\{|\xi_j| \leq N\}$ for $j = 1, 2$. Suppose also that the Fourier supports of u_1, u_2 are separated by at least $\geq cN$. Then for any $q > 2$,*

$$\|u_1 u_2\|_{L_{t,x}^q(I \times \mathbf{R})} \lesssim N^{1-3/q} \|u_1\|_{S_*^0(I \times \mathbf{R})} \|u_2\|_{S_*^0(I \times \mathbf{R})}, \quad (2.10)$$

where

$$\|u\|_{S_*^0(I \times \mathbf{R})} = \|u_0\|_{L^2(\mathbf{R})} + \|(i\partial_t + \Delta)u\|_{L_{t,x}^{6/5}(I \times \mathbf{R})}. \quad (2.11)$$

Proof: See [22].

To this end, let $q = 2 + \delta$, $\frac{1}{p} = \frac{1}{2} - \frac{1}{2+\delta}$.

In proving theorem 1.1, we will make use of a linear-nonlinear decomposition. See [19] for the linear-nonlinear decomposition for the defocusing, semilinear wave equation, [14] for the linear-nonlinear decomposition used for the three dimensional cubic defocusing nonlinear Schrödinger equation.

Theorem 2.3 *If*

$$\|\nabla Iu_0\|_{L^2(\mathbf{R})} \leq 1 \quad (2.12)$$

and for some $\epsilon > 0$ sufficiently small,

$$\|Iu\|_{L_t^{16/3}L_x^8(J \times \mathbf{R})} \leq \epsilon, \quad (2.13)$$

then

$$\|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \lesssim 1. \quad (2.14)$$

Moreover, the solution has the form

$$e^{it\Delta}u_0 + u^{nl}(t), \quad (2.15)$$

$$\|P_{>cN}\langle \nabla \rangle Iu^{nl}\|_{S^0(J \times \mathbf{R})} \lesssim \frac{1}{N^{1/2-}}. \quad (2.16)$$

Proof: The solution obeys the Duhamel formula,

$$Iu(t, x) = e^{it\Delta}u_0 + \int_0^t e^{i(t-\tau)\Delta}I(|u(\tau)|^4u(\tau))d\tau. \quad (2.17)$$

By the Strichartz estimates,

$$\|(1 - I)u\|_{L_t^{16/3}L_x^8(J \times \mathbf{R})} \lesssim \frac{1}{N} \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}, \quad (2.18)$$

$$\|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \lesssim \|\langle \nabla \rangle Iu_0\|_{L^2(\mathbf{R})} + \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \|u\|_{L_t^{16/3} L_x^8(J \times \mathbf{R})}^4 \quad (2.19)$$

$$\lesssim \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \lesssim \|\langle \nabla \rangle Iu_0\|_{L^2(\mathbf{R})} + \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} (\epsilon^4 + \frac{\|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^4}{N^4}). \quad (2.20)$$

Therefore, by the continuity method,

$$\|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \lesssim 1. \quad (2.21)$$

This takes care of (2.14). Next, we remark that this also proves

$$\|\langle \nabla \rangle I(|u|^4 u)\|_{L_{t,x}^{6/5}(J \times \mathbf{R})} \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^5 \lesssim 1. \quad (2.22)$$

To estimate the nonlinearity,

$$\nabla Iu^{nl}(t) = \int_0^t \nabla e^{i(t-\tau)\Delta} I(|u(\tau)|^4 u(\tau)) d\tau. \quad (2.23)$$

$$\begin{aligned} I(|u(\tau)|^4 u(\tau)) &= I(|u_{\leq \frac{cN}{10}}(\tau)|^4 u_{\leq \frac{cN}{10}}(\tau)) + I(O((u_{\leq \frac{cN}{100}}(\tau))^4 (u_{> \frac{cN}{10}}(\tau)))) \\ &\quad + I(O((u_{> \frac{cN}{10}}(\tau))(u_{> \frac{cN}{100}}(\tau))u(\tau)^3)). \end{aligned} \quad (2.24)$$

The first term, $I(|u_{\leq \frac{cN}{10}}(\tau)|^4 u_{\leq \frac{cN}{10}}(\tau))$ is supported on $|\xi| \leq \frac{cN}{2}$. To estimate the second term, use the bilinear estimates,

$$\begin{aligned} &\|\nabla I(O(u_{> \frac{cN}{10}}(u_{\leq \frac{cN}{100}})^4))\|_{L_t^1 L_x^2(J \times \mathbf{R})} \\ &\lesssim \|(\nabla Iu_{> \frac{cN}{10}})(u_{\leq \frac{cN}{100}})\|_{L_{t,x}^q} \|u_{\leq \frac{cN}{100}}\|_{L_{t,x}^p} \|u_{\leq \frac{cN}{100}}\|_{L_t^4 L_x^\infty}^2 \lesssim \frac{1}{N^{1/2-}}. \end{aligned}$$

Finally,

$$\begin{aligned} &\|\nabla I(O((u_{> \frac{cN}{10}}(\tau))(u_{> \frac{cN}{100}}(\tau))u(\tau)^3))\|_{L_t^1 L_x^2(J \times \mathbf{R})} \\ &\lesssim \|\nabla Iu\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|u_{> \frac{cN}{100}}\|_{L_t^{16/3} L_x^8(J \times \mathbf{R})} \|u\|_{L_t^{16/3} L_x^8(J \times \mathbf{R})}^3 \\ &\lesssim \frac{\epsilon^3}{N} \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^2 \lesssim \frac{1}{N}. \end{aligned}$$

The last estimate follows from

$$\begin{aligned}
\|u_{>\frac{cN}{100}}\|_{L_t^{16/3}L_x^8(J\times\mathbf{R})} &\lesssim \sum_{\frac{cN}{100}\leq N_j} \|P_{N_j}u\|_{L_t^{16/3}L_x^8(J\times\mathbf{R})} \\
&\lesssim \|\langle\nabla\rangle Iu\|_{S^0(J\times\mathbf{R})} \sum_{\frac{cN}{100}\leq N_j} \frac{1}{N_j^s N_j^{1-s}} \lesssim \frac{1}{N} \|\langle\nabla\rangle Iu\|_{S^0(J\times\mathbf{R})}.
\end{aligned} \tag{2.25}$$

□

3 Energy Increment

In this section we prove almost conservation of the modified energy $E(Iu(t))$.

Theorem 3.1 *Let*

$$E(Iu(t)) = \frac{1}{2} \int |\nabla Iu(t, x)|^2 dx + \frac{1}{6} \int |Iu(t, x)|^6 dx. \tag{3.1}$$

If J is an interval where a solution $u(t, x)$ of (1.1) exists, $\|u\|_{L_t^{16/3}L_x^8(J\times\mathbf{R})} \leq \epsilon$, $E(Iu_0) \leq 1$, then

$$\sup_{t_1, t_2 \in J} |E(Iu(t_1)) - E(Iu(t_2))| \lesssim \frac{1}{N^{3/2-}} \|P_{>cN} \nabla Iu\|_{L_t^4 L_x^\infty(J\times\mathbf{R})}^2 + \frac{1}{N^{2-}}, \tag{3.2}$$

where $c > 0$ is some small constant.

Proof:

$$\frac{d}{dt} E(Iu(t)) = \operatorname{Re} \int (\overline{Iu_t(t, x)}) [I(|u(t, x)|^4 u(t, x)) - |Iu(t, x)|^4 Iu(t, x)] dx. \tag{3.3}$$

Taking the Fourier transform, let $\Sigma = \{\xi_1 + \dots + \xi_6 = 0\}$, $d\xi$ is the Lebesgue measure on the hyperplane, using the fact that

$$Iu_t = iI\Delta u - iI(|u|^4 u), \tag{3.4}$$

$$\begin{aligned}
\frac{d}{dt} E(Iu(t)) &= -\operatorname{Re} \int_{\Sigma} (i|\xi_1|^2 \widehat{Iu}(t, \xi_1)) \left[1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2)m(\xi_3)\dots m(\xi_6)} \right] \\
&\quad \times \widehat{Iu}(t, \xi_2) \widehat{Iu}(t, \xi_3) \widehat{Iu}(t, \xi_4) \widehat{Iu}(t, \xi_5) \widehat{Iu}(t, \xi_6) d\xi
\end{aligned} \tag{3.5}$$

$$\begin{aligned}
& -Re \int_{\Sigma} (i\widehat{I(|u|^4 u)}(t, \xi_1)) [1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2)m(\xi_3) \cdots m(\xi_6)}] \\
& \quad \times \widehat{Iu}(t, \xi_2) \widehat{Iu}(t, \xi_3) \widehat{Iu}(t, \xi_4) \widehat{Iu}(t, \xi_5) \widehat{Iu}(t, \xi_6) d\xi.
\end{aligned} \tag{3.6}$$

We will estimate (3.5) and (3.6) separately by making a Littlewood-Paley decomposition and consider several cases separately. Without loss of generality let $N_2 \geq N_3 \geq N_4 \geq N_5 \geq N_6$.

The term (3.5):

When estimating this term, we will frequently use the bilinear estimate (2.10).

Case 1, $N_2 \ll N$: In this case, $m(\xi_i) \equiv 1$, so

$$1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \cdots m(\xi_6)} \equiv 0. \tag{3.7}$$

Case 2, $N_2 \gtrsim N \gg N_3$: By the fundamental theorem of calculus,

$$|1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \cdots m(\xi_6)}| \lesssim \frac{N_3}{N_2}. \tag{3.8}$$

Recall that $q = 2 + \delta$, $\frac{1}{p} = \frac{1}{2} - \frac{1}{q}$. $N_1 \sim N_2$, so

$$\begin{aligned}
& \sum_{N \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3 < N} \frac{N_1}{N_2^2} \|(P_{N_1} \nabla Iu)(P_{N_3} Iu)\|_{L_{t,x}^q} \\
& \quad \times \|(P_{N_2} \nabla Iu)(P_{N_4} Iu)\|_{L_{t,x}^q} \|P_{N_5} Iu\|_{L_{t,x}^p} \|P_{N_6} Iu\|_{L_{t,x}^p} \\
& \lesssim \sum_{N \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3 < N} \frac{N_1^{1/2+}}{N_2^{5/2-}} \frac{1}{\langle N_3 \rangle} \frac{1}{\langle N_4 \rangle \langle N_5 \rangle^{1/2-} \langle N_6 \rangle^{1/2-}} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

Case 3, $N_2 \geq N_3 \gtrsim N \gg N_4$: In this case, estimate the multiplier by

$$|1 - \frac{m(\xi_1)}{m(\xi_2)m(\xi_3)}| \lesssim \frac{m(\xi_1)}{m(\xi_2)m(\xi_3)}. \tag{3.9}$$

Consider three subcases separately.

Case 3(a), $N_3 \ll N_1 \sim N_2$:

$$\begin{aligned}
& \sum_{N \lesssim N_3 \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \ll N} \frac{1}{m(N_3)} \frac{N_1}{N_2} \|(P_{N_1} \nabla Iu)(P_{N_3} Iu)\|_{L_{t,x}^q} \\
& \quad \|(P_{N_2} \nabla Iu)(P_{N_4} Iu)\|_{L_{t,x}^q} \|P_{N_5} Iu\|_{L_{t,x}^p} \|P_{N_6} Iu\|_{L_{t,x}^p} \\
& \lesssim \sum_{N \lesssim N_3 \ll N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \ll N} \frac{N_1^{1/2}}{N_2^{3/2-}} \frac{1}{N_3^s N^{1-s}} \frac{1}{\langle N_4 \rangle \langle N_5 \rangle^{1/2-} \langle N_6 \rangle^{1/2-}} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

Case 3(b), $N_1 \ll N_2 \sim N_3$:

$$\begin{aligned}
& \sum_{N \lesssim N_2 \sim N_3} \sum_{N_6 \leq N_5 \leq N_4 \ll N, N_1 \ll N_2} \frac{m(N_1)}{m(N_2)m(N_3)} \frac{N_1}{N_2} \|(P_{N_2} \nabla Iu)(P_{N_1} \nabla Iu)\|_{L_{t,x}^q} \\
& \quad \|(P_{N_3} Iu)(P_{N_4} Iu)\|_{L_{t,x}^q} \|P_{N_5} Iu\|_{L_{t,x}^p} \|P_{N_6} Iu\|_{L_{t,x}^p} \\
& \lesssim \sum_{N \lesssim N_3 \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \ll N} \frac{1}{N_1^{1/2-} N_2^{1/2-}} \frac{1}{N_2^s N^{1-s}} \frac{N_1^s N^{1-s}}{N_3^s N^{1-s}} \\
& \quad \times \frac{1}{\langle N_4 \rangle \langle N_5 \rangle^{1/2-} \langle N_6 \rangle^{1/2-}} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

Case 3(c), $N_1 \sim N_2 \sim N_3$: In this case,

$$\begin{aligned}
& \frac{m(N_1)}{m(N_2)m(N_3)} \sim \frac{1}{m(N_3)}. \\
& \sum_{N \lesssim N_1 \sim N_2 \sim N_3} \sum_{N_6 \leq N_5 \leq N_4 \ll N} \frac{N_1}{N_2} \|(P_{N_1} \nabla Iu)(P_{N_4} Iu)\|_{L_{t,x}^q} \\
& \quad \|P_{N_2} \nabla Iu\|_{L_t^4 L_x^\infty} \|P_{N_3} Iu\|_{L_t^4 L_x^\infty} \|P_{N_5} Iu\|_{L_t^\infty L_x^2} \|P_{N_6} Iu\|_{L_{t,x}^p}
\end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{N \lesssim N_2 \sim N_3} \frac{1}{N_2^{1/2-} N^{1-s} N_3^s} \|P_{N_2} \nabla I u\|_{L_t^4 L_x^\infty} \|P_{N_3} \nabla I u\|_{L_t^4 L_x^\infty} \\
&\times \sum_{N_6 \leq N_5 \leq N_4 < N} \frac{1}{\langle N_4 \rangle \langle N_5 \rangle \langle N_6 \rangle^{1/2-}} \lesssim \frac{1}{N^{3/2-}} \|P_{>cN} I u\|_{L_t^4 L_x^\infty(J \times \mathbf{R})}^2.
\end{aligned}$$

Case 4, $N_2 \geq N_3 \geq N_4 \gtrsim N$:

Case 4(a), $N_1 \sim N_2$: In this case

$$\begin{aligned}
&|1 - \frac{m(\xi_1)}{m(\xi_2) \cdots m(\xi_6)}| \lesssim \frac{1}{m(\xi_3) \cdots m(\xi_6)}. \\
&\sum_{N \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3} \frac{N_1}{N_2} \|P_{N_1} \nabla I u\|_{L_{t,x}^6} \|P_{N_2} \nabla I u\|_{L_{t,x}^6} \\
&\quad \times \|P_{N_3} u\|_{L_{t,x}^6} \|P_{N_4} u\|_{L_{t,x}^6} \|P_{N_5} u\|_{L_{t,x}^6} \|P_{N_6} u\|_{L_{t,x}^6} \\
&\lesssim \sum_{N \lesssim N_1 \sim N_2} \sum_{N \lesssim N_4 \leq N_3} \sum_{N_6 \leq N_5 \leq N} \frac{N_1}{N_2} \|P_{N_1} \nabla I u\|_{L_{t,x}^6} \|P_{N_2} \nabla I u\|_{L_{t,x}^6} \\
&\quad \times \sum_{N \lesssim N_4 \leq N_3} \frac{1}{N_3^s N^{1-s}} \frac{1}{N_4^s N^{1-s}} \frac{1}{m(N_5) N_5} \frac{1}{m(N_6) N_6} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

Case 2, $N_1 << N_2$: In this case $N_2 \sim N_3$.

$$\begin{aligned}
&\sum_{N \lesssim N_2 \sim N_3} \sum_{N \lesssim N_4; N_6 \leq N_5 \leq N_4; N_1 << N_2} \frac{N_1}{N_2} \frac{m(N_1)}{m(N_2)} \|P_{N_1} \nabla I u\|_{L_{t,x}^6} \|P_{N_2} \nabla I u\|_{L_{t,x}^6} \\
&\quad \|P_{N_3} u\|_{L_{t,x}^6} \|P_{N_4} u\|_{L_{t,x}^6} \|P_{N_5} u\|_{L_{t,x}^6} \|P_{N_6} u\|_{L_{t,x}^6} \\
&\lesssim \sum_{N \lesssim N_2 \sim N_3} \sum_{N \lesssim N_4; N_6 \leq N_5 \leq N_4; N_1 << N_2} \frac{N_1}{N_2} \frac{m(N_1)}{m(N_2)} \frac{1}{N_3^s N^{1-s}} \frac{1}{N_4^s N^{1-s}} \\
&\quad \times \frac{1}{N_5 m(N_5)} \frac{1}{N_6 m(N_6)} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

This takes care of (3.5).

The term (3.6): Recall that this is the 10-linear term

$$Re \int_0^t \int \overline{iI(|u|^4 u)} [I(|u|^4 u) - |Iu|^4(Iu)] dx dt. \quad (3.10)$$

The term $I(|u|^4 u)$ poses a slight technical problem. Ideally, this term would be placed in $L_{t,x}^6$, and we would then repeat the analysis used in (3.5). However, in general this is not possible, so instead let $u_l = P_{\leq \frac{N}{30}} u$, $u_h = P_{> \frac{N}{30}} u$,

$$F(t, x) = \overline{I(|u|^4 u)} - I(\overline{O(u_h^4 u_l)}) - I(\overline{|u_h|^4 u_h}), \quad (3.11)$$

where $O(u_h^4 u_l)$ consists of those terms in $|u_l + u_h|^4(u_l + u_h)$ consisting of four u_h terms and one u_l term.

$$\begin{aligned} \|F(t, x)\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} &\lesssim \|\langle \nabla \rangle F(t, x)\|_{L_t^4 L_x^1(J \times \mathbf{R})} \\ &\lesssim \|\langle \nabla \rangle Iu\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|u_l\|_{L_{t,x}^\infty(J \times \mathbf{R})}^2 \|u\|_{L_t^\infty L_x^2(J \times \mathbf{R})}^2 \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^5. \end{aligned} \quad (3.12)$$

Now evaluate

$$\int_0^t \int_\Sigma \hat{F}(t, \xi_1) \left[1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \dots m(\xi_6)}\right] \widehat{Iu}(t, \xi_2) \widehat{Iu}(t, \xi_3) \widehat{Iu}(t, \xi_4) \widehat{Iu}(t, \xi_5) \widehat{Iu}(t, \xi_6), \quad (3.13)$$

via a Littlewood - Paley partition of unity and considering several subcases separately. Without loss of generality, let $N_2 \geq N_3 \geq \dots \geq N_6$.

Case 1, $N_2 \ll N$: Once again, $m(\xi_i) \equiv 1$, so

$$1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \dots m(\xi_6)} \equiv 0. \quad (3.14)$$

Case 2, $N_2 \gtrsim N \gg N_3$: In this case, apply the fundamental theorem of calculus,

$$\left|1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \dots m(\xi_6)}\right| \lesssim \frac{N_3}{N_2}. \quad (3.15)$$

$$\sum_{N \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3 \ll N} \frac{N_3}{N_2} \|P_{N_1} F\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_2} Iu\|_{L_t^4 L_x^\infty(J \times \mathbf{R})}$$

$$\begin{aligned}
& \times \|P_{N_3} Iu\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_4} Iu\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_5} Iu\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \|P_{N_6} Iu\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \\
& \lesssim \sum_{N \lesssim N_1 \sim N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3 < N} \frac{1}{N_2^2} \frac{1}{\langle N_4 \rangle} \frac{1}{\langle N_5 \rangle} \frac{1}{\langle N_6 \rangle} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

Case 3, $N_2 \geq N_3 \gtrsim N$ In this case make the crude estimate

$$|1 - \frac{m(\xi_2 + \dots + \xi_6)}{m(\xi_2) \cdots m(\xi_6)}| \lesssim \frac{1}{m(\xi_2) \cdots m(\xi_6)}. \quad (3.16)$$

$$\begin{aligned}
& \sum_{N_1 \lesssim N_2, N \lesssim N_3 \leq N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3} \|P_{N_1} F\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_2} u\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_3} u\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \\
& \times \|P_{N_4} u\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|P_{N_5} u\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \|P_{N_6} u\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \\
& \lesssim \sum_{N_1 \lesssim N_2, N \lesssim N_3 \leq N_2} \sum_{N_6 \leq N_5 \leq N_4 \leq N_3} \frac{1}{N_2^s N^{1-s}} \frac{1}{N_3^s N^{1-s}} \\
& \times \frac{1}{\langle N_4 \rangle m(N_4)} \frac{1}{\langle N_5 \rangle m(N_5)} \frac{1}{\langle N_6 \rangle m(N_6)} \lesssim \frac{1}{N^{2-}}.
\end{aligned}$$

It only remains to consider

$$\begin{aligned}
& Re \int_0^t \int (i \overline{I(|u_h|^4 u_h)}) [I(|u|^4 u) - |Iu|^4(Iu)] dx dt \\
& + Re \int_0^t \int i \overline{I(3|u_h|^4 u_l + 2|u_h|^2 u_h^2 \overline{u_l})} [I(|u|^4 u) - |Iu|^4(Iu)] dx dt.
\end{aligned} \quad (3.17)$$

To evaluate

$$\begin{aligned}
& Re \int_0^t \int (i \overline{I(|u_h|^4 u_h)}) I(|u|^4 u) dx dt \\
& + Re \int_0^t \int i \overline{I(3|u_h|^4 u_l + 2|u_h|^2 u_h^2 \overline{u_l})} I(|u|^4 u) dx dt,
\end{aligned} \quad (3.18)$$

it is necessary to take advantage of some cancellations.

$$\begin{aligned}
& Re \int_0^t \int (i \overline{I(|u_h|^4 u_h)}) I(|u_h|^4 u_h) dx dt \equiv 0. \\
& Re \int_0^t \int i \overline{I(3|u_h|^4 u_l + 2|u_h|^2 u_h^2 \overline{u_l})} I(|u_h|^4 u_h) dx dt
\end{aligned}$$

$$+Re \int_0^t \int i \overline{I(|u_h|^4 u_h)} (3|u_h|^4 u_l + 2|u_h|^2 u_h^2 \overline{u_l}) dx dt \equiv 0.$$

It remains to evaluate

$$\int_0^t \int I(|u_h|^4 u_h) I(O(u_l^2 u^3)) dx dt + \int_0^t \int I(O(u_h^4 u_l)) I(O(u_l u^4)) dx dt. \quad (3.19)$$

By (3.12),

$$\|I(u_l^2 u^3)\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^5,$$

so

$$\begin{aligned} \int_0^t \int I(|u_h|^4 u_h) I(O(u_l^2 u^3)) dx dt &\lesssim \|I(u_l^2 u^3)\|_{L_t^4 L_x^\infty(J \times \mathbf{R})} \|u_h\|_{L_t^4 L_x^\infty(J \times \mathbf{R})}^3 \|u_h\|_{L_t^\infty L_x^2(J \times \mathbf{R})}^2 \\ &\lesssim \frac{1}{N^5} \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^{10} \lesssim \frac{1}{N^5}. \end{aligned}$$

$$\int_0^t \int I(O(u_h^4 u_l)) I(O(u_l u^4)) dx dt \lesssim \|\langle \nabla \rangle I(u_h^4 u_l)\|_{L_t^2 L_x^1(J \times \mathbf{R})} \|I(u_l u^4)\|_{L_t^2 L_x^1(J \times \mathbf{R})}$$

$$\begin{aligned} &\lesssim \|\langle \nabla \rangle Iu\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \|u_l\|_{L_{t,x}^\infty(J \times \mathbf{R})} \|u_h\|_{L_{t,x}^6(J \times \mathbf{R})}^3 \|u_l\|_{L_{t,x}^\infty(J \times \mathbf{R})} \|u\|_{L_{t,x}^6(J \times \mathbf{R})}^3 \|u\|_{L_t^\infty L_x^2(J \times \mathbf{R})} \\ &\lesssim \frac{1}{N^3} \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})} \lesssim \frac{1}{N^3}. \end{aligned}$$

This takes care of (3.18). To finish estimating (3.17),

$$\begin{aligned} &Re \int_0^t \int (i \overline{I(|u_h|^4 u_h)}) (|Iu|^4 (Iu)) dx dt \\ &+ Re \int_0^t \int i \overline{I(3|u_h|^4 u_l + 2|u_h|^2 u_h^2 \overline{u_l})} (|Iu|^4 (Iu)) dx dt, \quad (3.20) \\ &\lesssim \|u_h\|_{L_{t,x}^6(J \times \mathbf{R})}^4 \|u\|_{L_{t,x}^6(J \times \mathbf{R})} \|Iu\|_{L_{t,x}^6(J \times \mathbf{R})} \|Iu\|_{L_{t,x}^\infty(J \times \mathbf{R})}^4 \lesssim \frac{1}{N^4}, \end{aligned}$$

and the proof of theorem 3.1 is complete. $\square$

4 Morawetz estimates

Theorem 4.1 *Let u be the solution to the nonlinear Schrödinger equation in one dimension,*

$$iu_t + \Delta u = |u|^4 u. \quad (4.1)$$

Then

$$\|Iu\|_{L_{t,x}^8([0,T] \times \mathbf{R})}^8 \lesssim \|Iu\|_{L_t^\infty \dot{H}_x^1([0,T] \times \mathbf{R})} \|u_0\|_{L^2(\mathbf{R})}^7 + \sum_{J_k} \frac{1}{N^{2-}} \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^{12}, \quad (4.2)$$

where $[0, T] = \cup_k J_k$.

Proof: We start with the case $I = 1$. We will use the method found in [13] and [6]. Let $\omega(t, z) : \mathbf{R} \times \mathbf{R}^4 \rightarrow \mathbf{C}$,

$$\omega(t, z) = u(t, x_1)u(t, x_2)u(t, x_3)u(t, x_4), \quad (4.3)$$

where $u(t, x)$ is a solution to (1.1). Then $\omega(t, z)$ obeys the equation

$$i\omega_t + \Delta_z \omega = \left(\sum_{i=1}^4 |u(t, x_i)|^4 \right) \omega(t, z) = \mathcal{N}. \quad (4.4)$$

Next, define the interaction Morawetz quantity,

$$M_a(t) = 2 \int_{\mathbf{R}^4} \partial_j a(z) \operatorname{Im}(\overline{\omega(t, z)} \partial_j \omega(t, z)) dz, \quad (4.5)$$

following the convention that repeated indices are summed. Let

$$T_{0j}(t, z) = 2 \operatorname{Im}(\overline{\omega(t, z)} \partial_j \omega(t, z)), \quad (4.6)$$

$$L_{jk}(t, z) = -\partial_{jk}(|\omega(t, z)|^2) + 4 \operatorname{Re}(\overline{\partial_j \omega} \partial_k \omega)(t, z), \quad (4.7)$$

$$\partial_t T_{0j} + \partial_k L_{jk} = 2\{\mathcal{N}, \omega\}_p^j. \quad (4.8)$$

$$\int_0^T \int_{\mathbf{R}^4} \partial_t \partial_j a(z) \operatorname{Im}(\overline{\omega(t, z)} \partial_j \omega(t, z)) dz dt \quad (4.9)$$

$$= \int_0^T \int_{\mathbf{R}^4} \partial_j a(z) \partial_{jkk}(|\omega(t, z)|^2) dz dt \quad (4.10)$$

$$-4 \int_0^T \int_{\mathbf{R}^4} \partial_j a(z) \partial_k \operatorname{Re}(\overline{\partial_j \omega} \partial_k \omega)(t, z) dz dt \quad (4.11)$$

$$+ 2 \int_0^T \int a_j(z) \{\mathcal{N}, \omega\}_p^j dz dt. \quad (4.12)$$

Now, evaluate each term separately. Make a change of variables, $y = Az$, where

$$A = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 \end{pmatrix} \quad (4.13)$$

is an orthonormal matrix with inverse

$$A^{-1} = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix}. \quad (4.14)$$

In the new variables, let $a(y) = (y_2^2 + y_3^2 + y_4^2)^{1/2}$,

$$-\Delta \Delta a(y) = 4\pi \delta(y_2, y_3, y_4).$$

$$A^{-1} \begin{pmatrix} y_1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} y_1 \\ y_1 \\ y_1 \\ y_1 \end{pmatrix}. \quad (4.15)$$

Therefore, integrating (4.10) by parts,

$$(4.10) = \int_0^T \int (-\Delta \Delta a(y)) |\omega(t, z)|^2 dz dt = 8\pi \int_0^T \int_{\mathbf{R}} |u(t, x)|^8 dx dt. \quad (4.16)$$

$\partial_{jk} a(z)$ is a positive semidefinite matrix, so integrating (4.11) by parts,

$$4 \int_0^T \int_{\mathbf{R}^4} (\partial_{jk} a(z)) \operatorname{Re}(\overline{\partial_j \omega} \partial_k \omega)(t, z) \geq 0. \quad (4.17)$$

Finally, for (4.12),

$$\{\mathcal{N}, \omega\}_p^j(t, z) = -2|\omega(t, z)|^2 \partial_j \left(\sum_{i=1}^4 |u(t, x_i)|^4 \right). \quad (4.18)$$

$$|\omega(t, z)|^2 \partial_j \left(\sum_{i=1}^4 |u(t, x_i)|^4 \right) = \frac{2}{3} \sum_{j=1}^4 \partial_j (|u(t, x_j)|^6 (|\prod_{i \neq j} |u(t, x_i)|^2)),$$

so integrating (4.12) by parts,

$$-2 \int_0^T \int a_j(z) \{\mathcal{N}, \omega\}_p^j(t, z) dz dt = \frac{4}{3} \int_0^T \int a_{jj}(z) |\omega(t, z)|^2 |u(t, x_j)|^4 dz dt \geq 0. \quad (4.19)$$

Therefore,

$$\begin{aligned} \int_0^T \int_{\mathbf{R}} |u(t, x)|^8 dx dt &\lesssim \left| \int_0^T \int \partial_t (a_j(z) \operatorname{Im} [\overline{\omega(t, z)} \partial_j \omega(t, z)]) dz dt \right| \\ &\lesssim \|u\|_{L_t^\infty \dot{H}^{1/2}([0, T] \times \mathbf{R})}^2 \|u_0\|_{L^2(\mathbf{R})}^6. \end{aligned} \quad (4.20)$$

Next, we prove an almost Morawetz estimate (see [5], [11], [15] for the two dimensional case; [5], [12], [6] for discussion of the one dimensional case).

If $u(t, x)$ solves (4.1), then $Iu(t, x)$ solves

$$iIu_t(t, x) + \Delta Iu(t, x) = I(|u(t, x)|^4 u(t, x)) = \mathcal{N}. \quad (4.21)$$

Split the nonlinearity into "good" and "bad" pieces, $\mathcal{N} = \mathcal{N}_g + \mathcal{N}_b$.

$$\mathcal{N}_g = \sum_{i=1}^4 |Iu(t, x_i)|^4 Iu(t, x_i) \prod_{j \neq i} Iu(t, x_j), \quad (4.22)$$

$$\mathcal{N}_b = \sum_{i=1}^4 [I(|u(t, x_i)|^4 u(t, x_i)) - |Iu(t, x_i)|^4 (Iu(t, x_i))] \prod_{j \neq i} Iu(t, x_j). \quad (4.23)$$

Let $\omega(t, z) = Iu(t, x_1)Iu(t, x_2)Iu(t, x_3)Iu(t, x_4)$, performing the same analysis will split

$$\int_0^T \int_{\mathbf{R}^4} \partial_t a_j(z) T_{0j}(t, z) dz dt,$$

into a sum of terms of the form (4.10), (4.11), and (4.12). If $\mathcal{N} = \mathcal{N}_g$, then the previous analysis would carry over identically. Indeed,

$$\int_0^T \int_{\mathbf{R}^4} (-\Delta \Delta a(z)) |\omega(t, z)|^2 dz dt = 8\pi \int_0^T \int |Iu(t, x)|^8 dx dt. \quad (4.24)$$

$$4 \int_0^T \int_{\mathbf{R}^4} (\partial_{jk} a(z)) \operatorname{Re}(\overline{\partial_j \omega} \partial_k \omega)(t, z) dz dt \geq 0. \quad (4.25)$$

$$2 \int_0^T \int a_j(z) \{\mathcal{N}_g, \omega\}_p^j(t, z) dz dt = \frac{4}{3} \int_0^T \int a_{jj}(z) |\omega(t, z)|^2 |Iu(t, x_j)|^4 dz \geq 0. \quad (4.26)$$

Therefore,

$$\int_0^T \int |u(t, x)|^8 dx dt \lesssim \|Iu\|_{L_t^\infty \dot{H}_x^1([0, T] \times \mathbf{R})} \|u_0\|_{L^2(\mathbf{R})}^7 + \left| \int_0^T \int a_j(z) \{\mathcal{N}_b, \omega\}_p^j(t, z) dz dt \right|. \quad (4.27)$$

To analyze the remainder $\mathcal{N}_b$, first consider a term of the form

$$\int_{J_k} \int a_j(z) \overline{\mathcal{N}_b(t, z)} \partial_j \omega(t, z) dz dt. \quad (4.28)$$

Recall (4.23), without loss of generality let $i = 1$ and estimate

$$\begin{aligned} & \int_{J_k} \int a_j(z) \overline{[I(|u(t, x_1)|^4 u(t, x_1)) - |Iu(t, x_1)|^4 Iu(t, x_1)] Iu(t, x_2) Iu(t, x_3) Iu(t, x_4)} \\ & \quad \times \partial_j (Iu(t, x_1) Iu(t, x_2) Iu(t, x_3) Iu(t, x_4)) dx dt. \end{aligned} \quad (4.29)$$

Because $a_j(z) \in L^\infty$,

$$(4.28) \lesssim \|I(|u(t, x)|^4 u(t, x)) - |Iu(t, x)|^4 Iu(t, x)\|_{L_t^1 L_x^2(J_k \times \mathbf{R})} \|\langle \nabla \rangle Iu\|_{L_t^\infty L_x^2(\mathbf{R})}^7. \quad (4.30)$$

To evaluate

$$I(|u(t, x)|^4 u(t, x)) - |Iu(t, x)|^4 Iu(t, x),$$

make a Littlewood - Paley partition of unity. Let

$$F(t, \xi) = \int_{\xi=\xi_1+\dots+\xi_5} \left[1 - \frac{m(\xi_1 + \dots + \xi_5)}{m(\xi_1) \dots m(\xi_5)}\right] \widehat{Iu}(t, \xi_1) \widehat{Iu}(t, \xi_2) \widehat{Iu}(t, \xi_3) \widehat{Iu}(t, \xi_4) \widehat{Iu}(t, \xi_5). \quad (4.31)$$

Without loss of generality, let $N_1 \geq N_2 \geq N_3 \geq N_4 \geq N_5$. Consider several cases separately.

Case 1: $N_1 \ll N$ In this case,

$$\left|1 - \frac{m(\xi_1 + \dots + \xi_5)}{m(\xi_1) \dots m(\xi_5)}\right| \equiv 0.$$

Case 2: $N_1 \gtrsim N \gg N_2$ In this case, by the fundamental theorem of calculus,

$$\left|1 - \frac{m(\xi_1 + \dots + \xi_5)}{m(\xi_1) \dots m(\xi_5)}\right| \lesssim \frac{N_2}{N_1}.$$

$$(4.28) \lesssim \sum_{N \lesssim N_1} \sum_{N_5 \leq N_4 \leq N_3 \leq N_2 \ll N} \frac{1}{N_1^2} \|P_{N_1} \langle \nabla \rangle Iu\|_{L_t^\infty L_x^2(J_k \times \mathbf{R})} \|P_{N_2} \langle \nabla \rangle Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})}$$

$$\times \|P_{N_3} \langle \nabla \rangle Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \|P_{N_4} \langle \nabla \rangle Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \|P_{N_5} \langle \nabla \rangle Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})}$$

$$\lesssim \frac{1}{N^{2-}} \|\langle \nabla \rangle Iu\|_{S^0(J \times \mathbf{R})}^5.$$

Case 3: $N_1 \geq N_2 \gtrsim N$ In this case, crudely estimate

$$\left|1 - \frac{m(\xi_1 + \dots + \xi_5)}{m(\xi_1) \dots m(\xi_5)}\right| \lesssim \frac{1}{m(N_1)m(N_2)m(N_3)m(N_4)m(N_5)}.$$

$$(4.28) \lesssim \sum_{N \lesssim N_2 \leq N_1} \|P_{N_1} u\|_{L_t^\infty L_x^2(J_k \times \mathbf{R})} \|P_{N_2} u\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})}$$

$$\times \sum_{N_5 \leq N_4 \leq N_3 \leq N_2} \|P_{N_3} u\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \|P_{N_4} u\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \|P_{N_5} u\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})}$$

$$\lesssim \sum_{N \lesssim N_2 \leq N_1} \frac{1}{N^{1-s} N_1^s} \frac{1}{N^{1-s} N_2^s}$$

$$\begin{aligned}
& \times \sum_{N_5 \leq N_4 \leq N_3} \frac{1}{m(N_3)N_3} \frac{1}{m(N_4)N_4} \frac{1}{m(N_5)N_5} \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^5 \\
& \lesssim \frac{1}{N^{2-}} \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^5.
\end{aligned}$$

For a term of the form

$$\int_{J_k} \int_{\mathbf{R}^4} a_j(z) (\partial_j \overline{\mathcal{N}_b(t, z)}) \omega(t, z) dz dt, \quad (4.32)$$

integrating by parts rewrites this term as a term of the form (4.28) plus a term of the form

$$\int_{J_k} \int_{\mathbf{R}^4} a_{jj}(z) (\overline{\mathcal{N}_b(t, z)}) \omega(t, z) dz dt. \quad (4.33)$$

By (4.23) and the estimates on (4.31),

$$\|\mathcal{N}_b(t, z)\|_{L_t^1 L_x^2(J_k \times \mathbf{R}^4)} \lesssim \frac{1}{N^{2-}} \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^8. \quad (4.34)$$

Δa does not lie in $L^\infty(\mathbf{R}^4)$, rather,

$$a_{jj}(y) \lesssim \frac{1}{(y_2^2 + y_3^2 + y_4^2)^{1/2}}. \quad (4.35)$$

Integrating

$$\frac{1}{(y_2^2 + y_3^2 + y_4^2)^{1/2}} Iu(t, x_1) Iu(t, x_2) Iu(t, x_3) Iu(t, x_4)$$

along the tube $y_2^2 + y_3^2 + y_4^2 \leq 1$, using (4.15),

$$\begin{aligned}
& \int_{-\infty}^{\infty} \int_{y_2^2 + y_3^2 + y_4^2 \leq 1} \frac{1}{y_2^2 + y_3^2 + y_4^2} |Iu(t, \frac{y_1 + y_2 + y_3 - y_4}{2})|^2 |Iu(t, \frac{y_1 + y_2 - y_3 + y_4}{2})|^2 \\
& \times |Iu(t, \frac{y_1 - y_2 + y_3 + y_4}{2})|^2 |Iu(t, \frac{y_1 - y_2 - y_3 - y_4}{2})|^2 dy_1 dy_2 dy_3 dy_4 \\
& \lesssim (\int_0^1 \frac{r^2}{r^2} dr) \|Iu\|_{L_t^\infty L_x^8(J_k \times \mathbf{R})}^8 \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^8,
\end{aligned}$$

so

$$\|a_{jj}(y) \omega(t, y)\|_{L_t^\infty L_y^2(J_k \times \mathbf{R}^4)} \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^4.$$

On the other hand, when $y_2^2 + y_3^2 + y_4^2 \geq 1$, $\partial_{jj}a(z)$ is bounded and

$$\|Iu(t, x_1)Iu(t, x_2)Iu(t, x_3)Iu(t, x_4)\|_{L_t^1 L_x^2(J_k \times \mathbf{R}^4)} \lesssim \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^4. \quad (4.36)$$

Since

$$\|I(|u(t, x_1)|^4 u(t, x_1)) - |Iu(t, x_1)|^4 Iu(t, x_1)\|_{L_t^1 L_x^2(J \times \mathbf{R})} \lesssim \frac{1}{N^{2-}} \|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^5,$$

theorem 4.1 is proved.

5 Proof of Theorem 1.1

Theorem 5.1 (1.1) is globally well-posed for $u_0 \in H^s(\mathbf{R})$, $s > \frac{8}{29}$.

Proof:

$$\begin{aligned} \int |\nabla Iu_0(x)|^2 dx &\lesssim N^{2(1-s)} \|u_0\|_{H^s(\mathbf{R})}. \\ \int |Iu_0(x)|^6 dx &\lesssim N^{2-6s} \|u_0\|_{H^s(\mathbf{R})}^6. \end{aligned} \quad (5.1)$$

If $u(t, x)$ solves (1.1) on $[0, T_0]$, then rescaling,

$$\frac{1}{\lambda^{1/2}} u\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right) \quad (5.2)$$

solves (1.1) on $[0, \lambda^2 T_0]$, we will call the rescaled solution $u_\lambda(t, x)$. Choose $\lambda \sim N^{(1-s)/s}$ so that $E(Iu_\lambda(0)) = 1/2$. Let

$$W = \{t : E(Iu_\lambda(t)) \leq \frac{9}{10}\}. \quad (5.3)$$

W is closed by the dominated convergence theorem and nonempty since $0 \in W$. To prove $W = [0, \lambda^2 T_0]$, it suffices to prove W is open in $[0, \lambda^2 T_0]$. Suppose $W = [0, T]$, then by continuity there exists $\delta > 0$ such that $E(Iu_\lambda(t)) \leq 1$ on $[0, T + \delta]$.

Lemma 5.2 When $s > 1/4$,

$$\|Iu_\lambda\|_{L_{t,x}^s([0, T+\delta] \times \mathbf{R})}^8 \leq \frac{3}{2} C_0, \quad (5.4)$$

for some $C_0(T_0)$.

Proof: Let $\tau = \sup\{T_* \in [0, T + \delta] : \|Iu_\lambda\|_{L_{t,x}^8([0, T_*] \times \mathbf{R})}^8 \leq \frac{3}{2}C_0\}$. If $\tau < T + \delta$, there exists $\delta' > 0$ such that

$$\|Iu_\lambda\|_{L_{t,x}^8([0, \tau + \delta'] \times \mathbf{R})}^8 \leq 2C_0. \quad (5.5)$$

Recall $E(Iu_\lambda(t)) \leq 1$ on $[0, \tau + \delta']$,

$$\|Iu_\lambda\|_{L_t^{16/3} L_x^8([0, \tau + \delta'] \times \mathbf{R})} \leq (\lambda^2 T_0)^{1/16} \|Iu_\lambda\|_{L_{t,x}^8([0, \tau + \delta'] \times \mathbf{R})} \leq (\lambda^2 T_0)^{1/16} (2C_0)^{1/8}. \quad (5.6)$$

Partition $[0, \tau + \delta']$ into

$$\frac{(\lambda^2 T_0)^{1/3} (2C_0)^{2/3}}{\epsilon^{16/3}}$$

subintervals such that

$$\|Iu_\lambda\|_{L_t^{16/3} L_x^8([0, \tau + \delta'] \times \mathbf{R})} \leq \epsilon \quad (5.7)$$

on each subinterval. Then apply the almost Morawetz estimate,

$$\begin{aligned} \|Iu_\lambda\|_{L_{t,x}^8([0, \tau + \delta'] \times \mathbf{R})}^8 &\leq C \|u_0\|_{L^2(\mathbf{R})}^7 \|Iu_\lambda\|_{L_t^\infty \dot{H}^1([0, \tau + \delta'] \times \mathbf{R})} + \sum_k \frac{\|\langle \nabla \rangle Iu\|_{S^0(J_k \times \mathbf{R})}^{12}}{N^{2-}} \\ &\leq C_0 + C \frac{(\lambda^2 T_0)^{1/3} (2C_0)^{2/3}}{\epsilon^{16/3} N^{2-}} \leq C_0 + C \frac{N^{\frac{2(1-s)}{3s}} T_0^{1/3} (2C_0)^{2/3}}{\epsilon^{16/3} N^{2-}} \leq \frac{3}{2} C_0, \end{aligned} \quad (5.8)$$

when N is sufficiently large, as long as $\frac{2}{3}(\frac{1-s}{s}) < 2$, or $s > 1/4$. This proves the lemma. $\square$

Returning to the theorem,

$$\begin{aligned} \|Iu_\lambda\|_{L_{t,x}^8([0, T + \delta] \times \mathbf{R})}^8 &\leq \frac{3}{2} C_0, \\ \|Iu_\lambda\|_{L_t^{16/3} L_x^8([0, T + \delta] \times \mathbf{R})} &\lesssim \lambda^{1/8} T_0^{1/16}. \end{aligned} \quad (5.9)$$

Partition $[0, T + \delta]$ into $\lesssim (\lambda^2 T_0)^{1/3}$ subintervals. We will call these the little intervals. Take the union of the first $N^{1/2-}$ little subintervals, and call this big interval

$$J_1 = \cup_{l=1}^{N^{1/2-}} J_{1,l}.$$

Take the union of the next $N^{1/2-}$ subintervals and call this big interval J_2 , and so on.

$$[0, T + \delta] = \bigcup_{k=1}^{\frac{(\lambda^2 T_0)^{1/3}}{N^{1/2-}}} J_k = \bigcup_{k=1}^{\frac{(\lambda^2 T_0)^{1/3}}{N^{1/2-}}} \bigcup_{l=1}^{N^{1/2-}} J_{k,l}. \quad (5.10)$$

Lemma 5.3 *Suppose $E(Iu(t)) \leq 1$ on J_k .*

$$\sup_{t_1, t_2 \in J_k} |E(Iu(t_1)) - E(Iu(t_2))| \lesssim \frac{1}{N^{5/4-}}. \quad (5.11)$$

Proof: By theorem (3.1),

$$\sup_{t_1, t_2 \in J_{k,l}} |E(Iu(t_1)) - E(Iu(t_2))| \lesssim \frac{1}{N^{3/2-}} \|P_{>cN} \nabla Iu\|_{L_t^4 L_x^\infty(J_{k,l} \times \mathbf{R})}^2 + \frac{1}{N^{2-}} \quad (5.12)$$

$$\sum_{l=1}^{N^{1/2}} \|P_{>cN} \nabla Iu\|_{L_t^4 L_x^\infty(J_{k,l} \times \mathbf{R})}^2 \lesssim N^{1/4} \|P_{>cN} \nabla Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})}^2.$$

Let $J_k = [a, b]$, $J_{k,m} = [a_m, b_m]$, $a_{m+1} = b_m$, $a_1 = a$, $b_{N^{1/2-}} = b$. On each little subinterval, perform the linear-nonlinear decomposition in theorem 2.3. The solution on $[a_m, b_m]$ is of the form

$$e^{i(t-a_m)\Delta} u(a_m) + u_m^{nl}(t).$$

By induction,

$$e^{i(t-a_m)\Delta} u(a_m) = e^{it\Delta} u_0 + \sum_{j=1}^{m-1} e^{i(t-a_j)\Delta} u_j^{nl}(b_j). \quad (5.13)$$

$$\|\nabla e^{i(t-a)\Delta} Iu(a)\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \lesssim 1. \quad (5.14)$$

$$\sum_{m=1}^{N^{1/2-}} \|\nabla e^{i(t-a)\Delta} Iu(b_m)^{nl}\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \lesssim 1. \quad (5.15)$$

$$\sum_{l=1}^{N^{1/2-}} \|\nabla Iu^{nl}(t)\|_{L_t^4 L_x^\infty(J_{k,l} \times \mathbf{R})}^4 \lesssim \frac{1}{N^{3/2-}}. \quad (5.16)$$

Therefore,

$$\|\nabla P_{>cN} Iu\|_{L_t^4 L_x^\infty(J_k \times \mathbf{R})} \lesssim 1.$$

Plugging this back in to (5.11),

$$\sum_{l=1}^{N^{1/2-}} \sup_{t_1, t_2 \in J_{k,l}} |E(Iu(t_1)) - E(Iu(t_2))| \lesssim \frac{1}{N^{5/4-}} + \frac{1}{N^{3/2-}},$$

which proves the lemma. $\square$

Returning to the theorem once again, if $(\lambda^2 T_0)^{1/3} \ll N^{7/4-}$, for $t \in [0, T + \delta]$, or if $s > \frac{8}{29}$,

$$E(Iu(t)) \leq \frac{1}{2} + C \frac{(\lambda^2 T_0)^{1/3}}{N^{7/4-}} = \frac{1}{2} + C \frac{N^{\frac{2(1-s)}{3s}} T_0^{1/3}}{N^{7/4-}} \leq \frac{9}{10}, \quad (5.17)$$

for sufficiently large N , proving the theorem. We have to take $N \sim T_0^{\frac{4s}{29s-8}}$. Since $\lambda \sim N^{\frac{1-s}{s}}$, and

$$\|u(t)\|_{H^s(\mathbf{R})} = \lambda^s \|u_\lambda(\frac{t}{\lambda^2})\|_{H^s(\mathbf{R})},$$

$$\sup_{t \in [0, T]} \|u(t)\|_{H^s(\mathbf{R})} \lesssim (1 + T)^{\frac{4(1-s)s}{29s-8}+}. \quad (5.18)$$

$\square$

References

- [1] J. Bourgain. Refinements of Strichartz' inequality and applications to 2D-NLS with critical nonlinearity. *International Mathematical Research Notices*, 5:253 – 283, 1998.
- [2] J. Bourgain. *Global Solutions of Nonlinear Schrödinger Equations*. American Mathematical Society Colloquium Publications, 1999.
- [3] T. Cazenave and F. Weissler. The Cauchy problem for the nonlinear Schrödinger Equation in H^1 . *Manuscripta Mathematica*, 61:477 – 494, 1988.
- [4] T. Cazenave and F. Weissler. The Cauchy problem for the nonlinear Schrödinger Equation in H^s . *Nonlinear Analysis*, 14:807 – 836, 1990.
- [5] J. Colliander, M. Grillakis, and N. Tzirakis. Improved interaction Morawetz inequalities for the cubic nonlinear Schrödinger equation on $\mathbf{R}^2$. *Int. Math. Res. Not. IMRN*, (23):90 – 119, 2007.
- [6] J. Colliander, J. Holmer, M. Visan, and X. Zhang. Global existence and scattering for rough solutions to generalized nonlinear Schrödinger equations on $\mathbb{R}$. *Commun. Pure Appl. Anal.*, 7(3):467–489, 2008.
- [7] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation. *Mathematical Research Letters*, 9:659 – 682, 2002.
- [8] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on $\mathbf{R}^3$. *Communications on Pure and Applied Mathematics*, 21:987 – 1014, 2004.
- [9] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Resonant decompositions and the I-method for cubic nonlinear Schrödinger equation on $\mathbf{R}^2$. *Discrete and Continuous Dynamical Systems A*, 21:665 – 686, 2007.
- [10] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in $\mathbb{R}^3$. *Ann. of Math. (2)*, 167(3):767–865, 2008.

- [11] J. Colliander and T. Roy. Bootstrapped Morawetz estimates and resonant decomposition for low regularity global solutions of cubic NLS on $\mathbf{R}^2$. *preprint, arXiv:0811.1803*.
- [12] D. De Silva, N. Pavlovic, G. Staffilani, and N. Tzirakis. Global well-posedness for the L^2 -critical nonlinear Schrödinger equation in higher dimensions. *to appear, Communications on Pure and Applied Analysis*.
- [13] D. De Silva, N. Pavlović, G. Staffilani, and N. Tzirakis. Global well-posedness and polynomial bounds for the defocusing L^2 -critical nonlinear Schrödinger equation in $\mathbb{R}$. *Comm. Partial Differential Equations*, 33(7-9):1395–1429, 2008.
- [14] B. Dodson. Global well-posedness for the defocusing, cubic, nonlinear Schrödinger equation when $n = 3$ via a linear-nonlinear decomposition. *arXiv:0910.2260*.
- [15] B. Dodson. Improved almost Morawetz estimates for the cubic nonlinear Schrödinger equation. *arXiv:0909.0757*.
- [16] M. Keel and T. Tao. Endpoint Strichartz estimates. *American Journal of Mathematics*, 120:955 – 980, 1998.
- [17] R. Killip, T. Tao, and M. Visan. The cubic nonlinear Schrödinger equation in two dimensions with radial data. *Journal of the European Mathematical Society*, to appear.
- [18] R. Killip, M. Visan, and X. Zhang. The mass-critical nonlinear Schrödinger equation with radial data in dimensions three and higher. *Anal. PDE*, 1(2):229–266, 2008.
- [19] T. Roy. Adapted linear-nonlinear decomposition and global well-posedness for solutions to the defocusing cubic wave equation on $\mathbb{R}^3$. *Discrete Contin. Dyn. Syst.*, 24(4):1307–1323, 2009.
- [20] C. Sogge. *Fourier Integrals in Classical Analysis*. Cambridge University Press, 1993.
- [21] E. Stein. *Harmonic Analysis: Real Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton University Press, 1993.
- [22] T. Tao. A sharp bilinear restrictions estimate for paraboloids. *Geom. Funct. Anal.*, 13(6):1359–1384, 2003.

- [23] T. Tao. *Nonlinear Dispersive Equations: Local and Global Analysis*. American Mathematical Society, 2006.
- [24] T. Tao, M. Visan, and X. Zhang. The nonlinear Schrödinger equation with combined power-type nonlinearities. *Comm. Partial Differential Equations*, 32(7-9):1281–1343, 2007.
- [25] M. Taylor. *Pseudodifferential Operators and Nonlinear PDE*. Birkhauser, 1991.
- [26] M. Taylor. *Partial Differential Equations*. Springer Verlag Inc., 1996.
- [27] Y. Tsutsumi. L^2 solutions for nonlinear Schrödinger equation and nonlinear groups. *Funkcional Ekvacioj*, 30:115 – 125, 1987.
- [28] M. Visan. The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions. *Duke Mathematical Journal*, 138:281 – 374, 2007.